Name:	Date:	Class:

## **Control Lab Procedures**

SAMPLE NAME: Control Reaction Scheme

Corn Starch	10 g
Water	60 ml
Vinegar	5 ml
Glycerin	5 ml
Phosphorescence Solid	0.6 g

Note: Make sure you add your qualitative and quantitative observations as you move through each procedure step.

Procedures	Observations
1. Pre-heat hot plate to 400°C	
2. Label the top and bottom of a clean petri dish with "control" and	
with your group number	
3. In a 1000 ml beaker add 10 g of cornstarch (using weigh paper	
and a balance)	
4. Add 5 ml of vinegar (using a 10 ml graduated cylinder) to the	
same beaker	
5. Add 5 ml of glycerin (using a 10 ml graduated cylinder) to the	
same beaker	
6. Add 60 mL of water (using a 100 mL graduated cylinder) to the	
same beaker	
7. Stir the mixture using a silicon spatula until the corn starch is	
dissolved and mixture is thoroughly combined	
8. Measure out 60 ml of the mixture (using a 100 ml graduated	
cylinder) and dispense into to a clean 250 ml beaker	
9. Add 0.6 g of the phosphorescence solid to the 60 ml mixture in	
the 250 ml beaker (from Step 8 using weigh paper and a balance)	
10. Stir the mixture using a clean silicon spatula until the	
phosphorescence solid is completely dissolved throughout.	
11. Heat the mixture in the 250 ml beaker using a hot plate	
preheated to 400°C	
12. Continuously stir the mixture with the silicone spatula while it is	
heating.	
13. Continue to heat and stir the mixture for 6 minutes and 30	
seconds.	
14. Transfer the heated mixture into the labeled petri dish using	
the silicon spatula and heat glove.	
15. Allow the sample to dry and harden overnight. Do not cover!	